

# Determination of $\beta$ -Globin Gene Cluster Haplotypes and Prevalence of $\alpha$ -Thalassemia in Sickle Cell Anemia Patients in Venezuela

A. Arends,<sup>1\*</sup> M. Alvarez,<sup>1</sup> D. Velázquez,<sup>1</sup> M. Bravo,<sup>2</sup> R. Salazar,<sup>2</sup> J.M. Guevara,<sup>1</sup> and O. Castillo<sup>1</sup>

<sup>1</sup>Laboratorio de Hemoglobinas Anormales, Servicio de Hematología "Dr. Tulio Arends" Hospital Universitario de Caracas, Caracas, Venezuela

<sup>2</sup>Universidad de Oriente, Núcleo Sucre, Cumaná, Venezuela

Sickle cell anemia and  $\alpha$ -thalassemia have a heterogeneous distribution in Venezuela with a high frequency in the coastal area (sea level) and few cases in the mountains. Most of our population is an ethnic admixture of Europeans (Spaniards colonists), Africans (slaves), and Amerindians. The purpose of our study was to determine the origin of the  $\beta^s$  globin haplotype, age and survival dependency, and the admixture among the different African groups in our population. The  $\alpha^{3,7}$  globin gene deletion status was also studied and found in a very high frequency. DNA from peripheral blood of 191 non-related patients (81 with HbS homozygous and 15 patients compound heterozygous for HbS, HbC, HbD with  $\beta$ -thalassemia, and 95 with sickle cell trait) were studied. The  $\beta^s$  chromosome was linked 51% to the Benin Haplotype, 29.5% with the CAR, 12.5% to the Senegal, and 2.5% to the Cameroon. We did not find any significant difference between the haplotype distribution among adults and children and among sickle cell patients and traits. Only 8.6% of the patients have homozygosity for the Benin haplotype. These results show a very high frequency of admixture in our African origin population. *Am. J. Hematol.* 64: 87–90, 2000. © 2000 Wiley-Liss, Inc.

**Key words:**  $\beta$ -globin haplotypes; sickle cell anemia;  $\alpha$ -thalassemia

## INTRODUCTION

The presence of Hb S has been known in Venezuela for 50 years, and it presents a variable distribution with a high frequency in the coastal area (sea level) and seldom cases in the mountains. Venezuela is located at the North of South America and it has a large coast in the Caribbean Sea. Most of our population is an ethnic admixture of Europeans (Spaniards colonists), Africans (slaves), Amerindians, and later on the Italian, Spaniards, and Portuguese immigrants [1]. The  $\beta^s$  globin has a variable presentation, in population surveys it has been found to have a frequency as high as 19%, in isolated coastal communities, and as low as 0.0% in the Andean regions [1–3].

In a small Venezuelan village, the  $\beta$  globin gene cluster haplotypes were studied in normal population. The Benin haplotype represented 20.9% among 12 haplotypes studied, which indicates an important contribution of an African ancestry in that population [4]. In a previous study of a smaller group of sickle cell anemia patients the most prevalent linkage of the  $\beta^s$  chromosome was to the Benin haplotype [5].

We wanted to determine the origin of the  $\beta^s$  globin gene in our sickle cell anemia (SCA) patients as well as the age dependency, as has been noted in Cuba and Guadeloupe [6,7]. In order to avoid distortions in the frequency of any specific group, we also studied the  $\beta$  gene cluster haplotypes among sickle cell traits. We explored the frequency of  $\alpha$ -thalassemia and if it had age dependency.

## SUBJECTS AND METHODS

The study was performed in 191 non-related patients proceeding from all over the country, under regular fol-

Contract grant sponsor: BID-CONICIT; Contract grant number: BTS-067 OC\VE.

\*Correspondence to: A. Arends, Apto 89758 El Hatillo, Caracas, Venezuela 1083. E-mail: aarends@ibm.net

Received for publication 5 November 1998; Accepted 1 December 1999

**TABLE I. Distribution of  $\beta^s$  Haplotypes in SCA and Sickie Trait (HbAS)\***

$\beta^s$ haplotypes	SCA		HbAS	
	No.	%	No.	%
Benin	90	50.8	49	51.6
CAR	57	32.2	24	25.3
Senegal	25	14.2	9	9.4
Cameroon	4	2.3	3	3.2
Atypical	1	0.5	10	10.5
Total	177		95	

\*No significant difference ( $\chi^2 = 5.3$ ,  $P = 0.1499$ ) was found between SCA and sickle trait patients.

low-up in the Hemoglobinopathy Clinic of the Hospital Universitario de Caracas and the Hospital Civil de Maracay; 81 of them with Hb S homozygous, 15 compound heterozygous with Hb C, Hb D, and  $\beta$ -thalassemia, and 95 with the sickle cell trait. All patients and their parents provided consent to participate in this study. Phenotypes were identified by hemoglobin electrophoresis in acetate and citrate agar, solubility tests, and family studies. Diagnosis of  $\beta$ -thalassemia was confirmed by mutation identification using ARMS-PCR [8]. DNA from peripheral blood leukocytes was prepared as previously described [9]. Seven fragments of the  $\beta$ -like globin gene cluster were amplified by PCR using the primers described by Sutton and co-workers [10]. They were digested with restriction endonuclease to identify the polymorphic sites *Xmn*I 5' to the  $G\gamma$  gene, *Hind*III in the IVSII of  $G\gamma$  and  $A\gamma$  gene, *Hinc*II 3' and inside the  $\phi\beta$  gene, and *Hinf*I, *Hpa*I, respectively, at 5' and 3' of the  $\beta$  gene. Using these seven polymorphisms, the chromosomes were grouped in 5 haplotypes: CAR (−+−+−+), Benin (−−−+−−), Senegal (+−+−+−+), Cameroon (−+−+−+), and Atypical haplotypes (all other combinations). The  $-\alpha^{3.7}$  thalassemia cases were detected by GAP-PCR [11].

## RESULTS

The haplotypes of  $\beta^s$  chromosomes determined in 191 patients (272 chromosome) is shown in Table I. Distribution of the 177 chromosomes from the SCA patients studied was as follows: Benin haplotypes (Ben) 50.8%, followed by Bantu (CAR) 32.2%, Senegal (Sen) 14.1%, Cameroon (Cam) 2.3%, and atypical haplotypes (Aty) 0.5%. In the 95 sickle cell trait patients the distribution of the  $\beta^s$  gene was as follows: Ben 51.6%, followed by CAR 25.3%, Sen 9.4%, Cam 3.2%, and Aty 10.5%. The frequency distribution of the haplotypes was compared between SCA and HbAS, and no statistical difference ( $\chi^2 = 5.3$ ,  $P = 0.1499$ ) was noted among both groups. This was done in order to avoid a variable due to early death in a specific group. When age was considered, we found

**TABLE II. Distribution by Age of  $\beta^s$  Globin Gene Haplotypes in SCA Patients\***

$\beta^s$ haplotypes	Adults ( $n = 43$ )		Children ( $n = 53$ )	
	No.	%	No.	%
Benin	45	52.9	45	48.9
CAR	25	29.4	32	34.7
Senegal	12	14.2	13	14.2
Cameroon	3	3.5	1	1.1
Atypical			1	1.1
Total	85		92	

\*No statistically significant difference ( $\chi^2 = 1.3$ ,  $P = 0.5226$ ) was found between age group.

no statistical difference in the different age groups ( $\chi^2 = 1.3$ ,  $P = 0.5226$ ; Table II). We found a very high frequency of mixed haplotypes 82%, and only 8.6% were homozygous for Benin haplotype.

The  $-\alpha^{3.7}$   $\beta$  thalassemia mutation was the most prevalent among the compound heterozygous Hb S- $\beta$  thalassemia patients.

In 148 patients we found 92 (62.2%) with both  $\alpha$  genes normal, 48 (32.4%) heterozygous for  $-\alpha^{3.7}$  thalassemia, and 8 (5.4%) were homozygous for the same deletion. When age distribution was studied among the patients, with one or both  $\alpha$  genes deleted, no difference between age groups was found (Table III). There is a very high frequency of  $-\alpha^{3.7}$  thalassemia (0.27) in the 296 patients studied.

## DISCUSSION

The  $\beta^s$  mutation may be associated with at least four different major  $\beta^s$  haplotypes, named according to the geographic areas where it predominates: the Benin, the Senegal, the Central African Republic, and the Asian types [12,13]. Historically it is known that some slave vessels came straight from Africa to Venezuela. Afterward slaves were brought over as a result of licit commercial transactions from the Antilles for agriculture and mining works. Runaway slaves from Curacao and Trinidad arrived in Venezuela surreptitiously remaining free in the jungle, they were very aggressive and called "Black Cimarrons." Later they formed the population we find currently in the mountains. From an anthropologic point of view it is certain that the African population arriving in Venezuela possibly constituted a heterogeneous group which proceeded apparently from well-delimited African regions, Sudan and Bantu [14]. The studies of the  $\beta^s$  gene cluster are useful in the definition of the origin of the population brought by the slave trades. The incidence of  $\beta^s$  haplotypes in our country could be different according to the region studied as we know the frequency of HbS is variable and spread all over the country, been stronger in populations stemming from around the sites where the African first resided.

**TABLE III. α Thalassemia Frequency in Patients from Cuba, Guadeloupe, and Venezuela According to Age Group\***

Age	Cuba		Guadeloupe		Venezuela	
	Frequency	No.	Frequency	No.	Frequency	No.
0–20	0.14	74	0.18	169	0.30	64
21–50	0.19	144	0.2	110	0.23	32
>50	0.34	16	0.2	10	0.37	4
Total		234		289		100

\*In the Guadeloupean and Venezuelan groups no statistically significant difference was found. In the Cuban group there was a significant difference between the age groups.

In our study, we found the Benin β<sup>s</sup> haplotype as the most prevalent form, same as was found in Cuba, Guadeloupe, United States, and Jamaica [5,6,15,16], followed by the CAR haplotype, the most prevalent one in Brazil [17,18] (Table IV). Our findings were to be expected on the basis of African ancestry according to Curtin et al. [19], Nagel [16], and Pagnier [12]. These slaves had come from Central West African ports. The African Negroes brought to Brazil, monopoly of the Portuguese, were mostly from Angola, Congo, and Mozambique [19] where the CAR haplotype predominates. We found a very high frequency of mixed β<sup>s</sup> haplotypes in 82% of our patients, as was expected due to the high ethnic admixture.

The prevalence of −α<sup>3.7</sup> thalassemia in 37.8% of the subjects, with one or both chromosomes deleted, is one of the highest reported. This finding agrees with a previous report of a smaller group of patients [20]. No statistically significant difference was observed between children and adult with −α<sup>3.7</sup> thalassemia.

The studies by Nagel [21], Kusolik [22], Powars [23], Steinberg [24,25], and Chang [26] have demonstrated that the different β<sup>s</sup> gene cluster polymorphism modulates the clinical severity of the disease. The haplotypes and gender interact to modify the hematological parameters. They also may explain the phenotypic heterogeneity among the patients. As most of our patients of the Ben/CAR mixed haplotype, they should have the same clinical course, “benign/worst,” which is not what we see in our clinical practice. Even siblings with the same mixed haplotypes, with −α<sup>3.7</sup> thalassemia, the same environment, and socioeconomic conditions have different clinical behavior. This is one of the reasons for our belief that the problem has not been solved and that there must be other genetic variables modulating the clinical expression of our patients.

## ACKNOWLEDGMENTS

This work is dedicated to Dr. Tulio Arends, who initiated and encouraged us to follow his dream. We are very grateful to Maira Avila, Ana Chiarello, and Olivia

**TABLE IV. Distribution of β<sup>s</sup> Haplotypes in American Countries\***

β <sup>s</sup> haplotypes	Brazil	Cuba	Guadeloupe	Jamaica	EEUU	Venezuela <sup>a</sup>
Benin	25	51	74	72	62	51
CAR	73	41	11	17	17	32
Senegal	1	8	6	10	10	14

\*Data from Zago [16], Muñiz [5], Kéclard [6], Antonarakis [14], and Nagel [19].

<sup>a</sup>Report from present study.

Loreto for their technical assistance, and to Dr. Ana Cecilia Contreras and Dr. Joaquin Inaty for sharing some of their patients for this study. We thank Trina Arends for her advice and critical review of the manuscript.

## REFERENCES

1. Arends T, Gallango ML, Muller A, Gonzalez MM, Perez BO. Tapipa: A Negroid Venezuelan isolate. In: Meyer RJ, Otten CM, Ab del Hameed F, editors. *Evolutionary Models and Studies in Human Diversity*. The Hague: Mouton; 1978.
2. Arends T. Epidemiology of hemoglobin variants in Venezuela. In: Arends T, Bemsy G, Nagel R, editors. *Genetical, Functional and Physical Studies of Hemoglobin*. Basel: S. Karger; 1971.
3. Arends T. Estado actual del estudio de las hemoglobinas anormales en Venezuela. *Sangre* 1963;8:1–14.
4. Castro de Guerra D, Hutz MH, Bortolini MC, Mauro SF. β-Globin gene cluster haplotypes in an admixed Venezuelan Population. *Am J Hum Biol* 1997;9:323–327.
5. Arends A, Alvarez M, Velazquez D, Chiarello A, Salazar R, Guevara JM, Castillo O. Determination of β-globin gene cluster haplotypes and prevalence of thalassemia in sickle cell anemia patients in Venezuela. *Br J Haematol* 1998;102(1):48B (abstract).
6. Muñiz A, Corral L, Alaez C, Svarch E, Espinosa E, Carbonell N, Di Leo R, Felicetti L, Nagel RL, Martinez G. Sickle cell anemia and β gene cluster haplotypes in Cuba. *Am J Hematol* 1995;49(2):163–164.
7. Kéclard L, Romana M, Lavocat E, Saint-Martin C, Berchel C, Merault G. Sickle cell disorder, β-globin gene cluster haplotypes and α-thalassemia in neonates and adults from Guadeloupe. *Am J Hematol* 1997; 55:24–27.
8. Old JM, Varawalla NY, Weatherall DJ. Rapid detection and prenatal diagnosis of β thalassaemia: Studies in Indian and Cypriot populations in the UK. *Lancet* 1990;336:834–837.
9. Poncz M, Solwiejczyk D, Harpel B, Mory Y, Schwartz E, Surrey S. Construction of human genes libraries from small amount of peripheral blood: Analysis of β-like globin genes. *Hemoglobin* 1982;6:27–36.
10. Sutton M, Boushassira EE, Nagel RL. Polymerase chain reaction amplification applied to the determination of β-thal globin gene cluster haplotypes. *Am J Hematol* 1989;32:66–69.
11. Baysal E, Huisman THJ. Detection of common deletion α-thalassemia-2 determinants by PCR. *Am J Hematol* 1994;46:208–213.
12. Pagnier J, Mears JG, Dunda BO, Schaefer RKE, Beldjord C, Nagel RL, Labie D. Evidence of the multicenter origin of the hemoglobin S gene in Africa. *Proc Natl Acad Sci USA* 1984;81:1771–1773.
13. Kulozik AE, Wainscoat JS, Serjeant GR, Kar BC, Alejandra AB, Essan GJF, Falusi AG, Haque SK, Hilali AM, Kate S, Renasinghe WAEP, Weatherall DJ. Geographical survey of β-globin gene haplotypes: Evidence for an independent Asian origin of the sickle cell mutation. *Am J Hum Genet* 1986;39:239–244.

14. Acosta Saignes M, editor. *Vida de los esclavos negros en Venezuela*. Caracas Hesperides 1967.
15. Antonarakis SE, Boehm CD, Serjeant GR, Theisen CE, Dover GJ, Kazazian HH. Origin of the  $\beta^s$  globin genes in Blacks: The contribution of recurrent mutation or gene conversion or both. *Proc Natl Acad Sci USA* 1984;81:853–856.
16. Nagel RL. The origin of the hemoglobin S gene: Clinical, genetic and anthropological consequences. *Einstein Q J Med* 1984;2:53–62.
17. Zago MA, Melo SEJ, Clegg JB, Guerreiro JF, Martinson JJ, Norwich J, Figueiredo MS.  $\alpha$ -Globin gene haplotypes in South American Indians. *Hum Biol* 1995;67:535–546.
18. Costa FF, Arruda VR, Goncalves MG, Miranda SR, Carvalho MH, Sonati MF, Saad SO, Gesteira F, Fernandes D, Nascimento ML.  $\beta^s$  gene cluster haplotypes in sickle cell anemia patients from two regions of Brazil [letter]. *Am J Hematol* 1994;45:96–97.
19. Curtin PD. *The Atlantic Slave Trade: A Census*. Milwaukee: University of Wisconsin Press; 1969.
20. Arends A, Velazquez D, Avila M, Amesty C, Chiarello A, Guevara JM, Cavazza M, Salazar R, Castillo O. Detection of deletional  $\alpha$ -thalassemia-2 determinants in the Venezuelan population by PCR. *Int. J Hematol* 1996;64(1):S81 (abstract).
21. Nagel RL. Severity, pathobiology, epistatic effects and genetic markers in sickle cell anemia. *Semin Hematol* 1991;28:180–201.
22. Kulozik AE, Kar BC, Satapathy RK, Serjeant BE, Weatherall DJ. Fetal hemoglobin levels and  $\beta^s$  globin haplotypes in an Indian population with sickle cell diseases. *Blood* 1987;69:1742–1746.
23. Powars DR, Chan L, Schroeder WA.  $\beta^s$  gene cluster haplotypes in sickle cell anemia: Clinical implications. *Am J Pediatr Hematol Oncol* Fall 1990;12:367–374.
24. Steinberg MH, Hsu H, Nagel RL, Milner PF, Adams JG, Benjamin L, Fryd S, Gillette P, Gilman J, Josifovska O, Hellman-Erlingsson S, Safaya S, Huey L, Rieder RF. Gender and haplotype effects upon hematological manifestations of adult sickle cell anemia. *Am J Hematol* 1995;48:175–181.
25. Steinberg MH, Lu ZH, Nagel RL, Venkataramani S, Milner PF, Huey L, Safaya S, Rieder RF. Hematological effects of atypical and cameroonian  $\beta$  globin gene haplotypes in adult sickle cell anemia. *Am J Hematol* 1998;59:121–126.
26. Chang YC, Smith KD, Moore RD, Searjant GR, Dover GJ. An analysis of fetal hemoglobin variation in sickle cell disease: The relative contributions of the X-linked factor,  $\beta$  globin haplotypes,  $\alpha$  globin gene number, gender, and age. *Blood* 1995;85:1111–1117.